

**21st International Conference on
Harmonisation within Atmospheric Dispersion Modelling for Regulatory Purposes
27-30 September 2022, Aveiro, Portugal**

**EVALUATION OF THE EFFECTS OF THE NATIONAL EMISSION REDUCTION
STRATEGIES FOR YEARS 2020-2029 AND AFTER 2030 ON THE SULPHUR AND
NITROGEN SURFACE CONCENTRATIONS ON THE TERRITORY OF BULGARIA**

Ivelina Georgieva¹, Georgi Gadzhev¹, Kostadin Ganev¹ and Vladimir Ivanov¹

¹ National Institute of Geophysics, Geodesy and Geography, Bulgarian Academy of Sciences

Abstract: Complying the EU Directive 2016/2284, Bulgaria developed national emission reduction strategies for years 2020-2029 and after 2030. Evaluation of the effects of these strategies on the Sulphur and Nitrogen surface concentrations on the territory of Bulgaria is the objective of the present study.

The studies are performed applying computer simulations. The simulations are performed with the US EPA Models-3 System: Meteorological model WRF; Emission model SMOKE; Atmosphere Composition Model CMAQ for the period 2008 – 2014. The provided model simulations are with horizontal resolution 9 kilometers for the region of Bulgaria. The NCEP Global Analysis Data meteorological background with 1°x1° resolution is used as a meteorological background. The models nesting capabilities were applied to downscale the simulations to 9 km resolution for Bulgaria.

Five emission scenarios are considered: 2005 emissions (reference period), 2020-2029 emissions projected with existing measures (WEM) and with additional measures (WAM), projected after 2030 WEM and WAM emissions. The comparison of the concentrations, simulated with the different scenarios makes it possible to evaluate the effect of the national emission reduction strategies.

Key words: *emission reduction, surface concentrations, computer simulations, emission scenarios.*

INTRODUCTION

The present paper presents part of the results obtained in the frame of an extensive study of the effects of different emission reduction scenarios on the air quality in Bulgaria. The set of models applied for atmospheric composition simulations is the same used in the operational Bulgarian Chemical Weather Forecast and Information System. Thus, the obtained results are fully compatible with the operational chemical weather forecast. The models are also widely used in air pollution modelling, so the obtained computer simulation results are in harmony with evaluations made for other regions. Air pollution forces many countries to take action to mitigate its adverse effects on human health and ecosystems. There is already a large amount of direct and indirect data connected to the air quality from the different surface and satellite-based observing systems. However, we need to track the various processes involved in the composition, transport, and transformation of the air pollutant species, which will help us better understand their distribution at different spatial/temporal scales. These tasks are performed by air quality modeling systems, as we use US EPA Models 3 System for modeling the air quality in our country - Bulgaria.

The main goal of this work is to make a comparison between Nitrogen (NO₂) and Sulfur (SO₂) dioxide surface concentrations, simulated with the different emission reduction scenarios (with and without measures) and the reference period (with emission for 2005), in Bulgaria.

METHODOLOGY

Based on 3D modelling tools, an extensive database was created and used for different studies of atmospheric composition. The simulations were based on the US EPA Model-3 system. - WRF v.3.2.1 (Shamarock et al. 2007), (UCAR/NCAR) - Weather Research and Forecasting Model, used as meteorological pre-processor; CMAQ v.4.6 - Community Multi-Scale Air Quality model (Byun et al., 1998), (Byun and Ching, 1999), (CMAS) the Chemical Transport Model (CTM), and SMOKE - the Sparse Matrix Operator Kernel Emissions Modelling System (CEP, 2003) the emission pre-processor of Models-3 system. This group of models is called Models-3 and is a typical example of the models created by a large

number of teams in the implementation of various projects, funded mainly by the United States Environment Agency (US EPA) - so-called “Community models”. The models in the system were adapted and validated for Bulgaria. This gave the opportunity to conduct extensive studies on a fully competitive modern level of the climate of atmospheric composition in the country. Previous results from air pollution modeling for Bulgaria are published in many papers (Gadzhev et al., 2013a, 2013b, 2013c), (Georgieva E. et al., 2017, 2019), (Georgieva I., 2014), (Syrovak et al., 2012, 2014, 2015, 2019).

The studies are performed by applying computer simulations for a 7-year period (2008 – 2014). The provided model simulations are with a horizontal resolution of 9 kilometers for the territory of Bulgaria. The large-scale (background) meteorological data used were from the ‘NCEP Global Analysis Data’ with a horizontal resolution of 1°x1°. Using the ‘nesting’ capabilities of the models, a resolution of 9 km was achieved for the territory of Bulgaria (Fig.1). The resolution of the mother domain (Europe) is 81 km, and two other domains are nested in it and in each other – Balkan Peninsula (27 km resolution) and Bulgaria (9 km), as shown in Figure 1.

TNO inventory (Denier van der Gon et al., 2010) is exploited for the territories outside Bulgaria in the mother domain. For the Bulgarian domain, the National inventory as provided by Bulgarian Executive Environmental Agency is used.




Figure 1. Model domains: Downscaling from domain Europe (81 km) to domain Bulgaria (09 km) and pointed location of some industrial zones in Bulgaria – 1 - TPP Bobovdol, 2 – Sofia city, 3 – Plovdiv city, 4 - TPP Maritsa Iztok, 5 – Burgas city, 6 - Devnya Industrial Area.

Every EU member state (According to the legislation) has to report the set of emission projections scenarios (EEA Technical report No 4/2015): projections scenario 'with existing measures' (WEM) means projections of anthropogenic Greenhouse Gas (GHG) or air pollutant emissions by sources that encompass the effects of currently implemented or adopted policies and measures; projections scenario 'with additional measures' (WAM) means projections of anthropogenic GHG or air pollutant emissions by sources that encompass the effects of policies and measures which have been adopted and implemented, as well as planned policies that are judged to have a realistic chance to be adopted and implemented in the future;

In the paper, four emission scenarios are considered for two periods, 2020-2029 and after 2030, with existing measures (WEM) and with additional measures (WAM), and the results are compared with the reference period (Ref 2005). It should be clear that only the Bulgarian emissions for 2020-2029 and 2030 have been modified according to the forecast scenarios.

The considerations and conclusions in the paper are based on simulated SO₂ and NO₂ surface concentrations on the territory of Bulgaria. Comparing the concentrations simulated with the different scenarios makes it possible to evaluate the effect of the national emission reduction strategies (EU Directive 2016/2284).

RESULTS

The Normalized Mean Bias noted as NMB (1) was used for comparison of the effect of different scenarios. The notions in these equations are *i* - *i*th value, *S* - the scenario output (WEM/WAM) for the selected period (2020-2029, 2030), and *R* - the output for the Reference period (2005). All of these scenarios are presented separately for the SO₂ and NO₂ surface concentrations and comparison with the reference period (2005) together with NMB in the following figures (Figure 2, Figure 3).

$$NMB = \frac{\sum_i S - \sum_i R}{\sum_i R} * 100[\%] \quad (1)$$

Figure 2 presents the results from numerical experiments about NO₂ surface concentrations for different scenarios and the comparison with the results from the reference period 2005 and calculated NMB. According to these results, we can say that not only the measures for reductions of emissions worked, but also clearly see which scenario gives better results.




Figure 2. (a) Annually averaged NO₂ surface concentrations [μm^3] for Ref 2005, WEM 2029, WAM 2029, WEM 2030, WAM 2030 emissions scenarios and (b) NMB of the NO₂ surface concentrations [%] for WEM 2029, WAM 2029, WEM 2030, WAM 2030 emissions scenarios compared to the Ref 2005.

The results from all scenarios have a difference with a reduction tendency according to the reference period 2005. It is very well displayed for all plots and we can see that the values for some areas are smaller than those of Ref 2005 – 1.Sofia, 3.Plovdiv and 5.Burgas city, also 6.Devnya industrial area.

It can be stated that the scenario WAM 2030 shows significant reduction of the surface concentrations for the whole territory of the country. The NO₂ surface concentrations have a reduction for the WEM/WAM scenarios according to the Reference 2005, but on Figure 2(a) we cannot say which one of the scenarios is “working” better. All the statements are confirmed by the plots, where the NMB is calculated (b). The scale is up to 0, and all the biases are with a negative sign. That suggests that all the scenarios produce smaller concentrations compared to the reference period and emissions for 2005. All the measures show a positive effect of reduction of emissions and lead to a reduction of NO₂ surface concentrations. It can be clearly seen about 25 % reduction over the whole domain and about 40% for some areas in the southwest part of the country. The WAM scenarios show bigger reduction for both selected periods, but for WAM 2030 the effect is most strongly manifested.




Figure 3. (a) Annually averaged SO₂ surface concentrations [μm^3] for Ref 2005, WEM 2029, WAM 2029, WEM 2030, WAM 2030 emissions scenarios and (b) NMB of the SO₂ surface concentrations [%] for WEM 2029, WAM 2029, WEM 2030, WAM 2030 emissions scenarios compare to the Ref 2005.

Figure 3 presents the results of the SO₂ for the different scenarios. Here, the effects of the emission reduction on SO₂ surface concentrations are very well expressed, and the differences with Ref 2005 are obvious. Again, all the scenarios show better results according to the reference period. The SO₂ plots on Figure 3(a) also shows that the scenarios of reduction of the emission lead to a decrease in the SO₂ concentrations. For all the results, we can say that more stringent measures lead to more effective results, where as in this study this is about the WAM scenario. The better results show the WAM 2030 scenario compared with the

reference period and also compare with other scenarios. The results from plots on Figure 3(b) about the bias of the SO₂ concentrations, as we mention above, also have a negative sign. Here, the percentage of reduction is very well expressed and shows very good results of taking measures. The reduction reaches 50% almost over the whole territory of the country and for all scenarios. On the plots, we can also see a reduction of 70% for some areas in the southwest part where are situated 4. TTP Maritsa Iztok and 3. Plovdiv city.

CONCLUSIONS

Complying with the EU Directive 2016/2284, Bulgaria developed national emission reduction strategies for 2020-2029 and after 2030. Evaluation of the effects of these strategies on the SO₂ and NO₂ surface concentrations on the territory of Bulgaria is demonstrated by some comparison of the concentrations, simulated with the different scenarios. This comparison makes it possible to evaluate the effect of the national emission reduction strategies: The NMB is used, and according to the results, we can conclude that the measures for reduction of emissions have good results and determine which scenario is better: The numerical experiments about NO₂ surface concentrations for different scenarios show:

- the results from all scenarios have a difference with reduction tendency in comparison with the reference period 2005;
- the NO₂ surface concentrations are reduced for all the WEM/WAM scenarios according to the Reference 2005, but the scenario WAM 2030 shows better results for the whole territory of Bulgaria;
- all the biases are with a negative sign and show that all the scenarios have smaller concentrations compared with the reference period and emissions for 2005;
- all the measures show a positive effect of reduction of emissions and lead to a reduction of NO₂ surface concentrations;
- the reduction over the whole domain is about 25 % and about 40% for some areas in the southwest part of the country;
- the WAM scenarios show a more significant reduction for both selected periods, and also, the WAM 2030 is clearly the one with the most sound effect.

The numerical experiments about SO₂ surface concentrations for different scenarios show:

- the effects of the emission reduction on SO₂ surface concentrations are even better expressed, and the differences with Reference 2005 are obvious;
- all the scenarios show better results according to the reference period, which shows that all the scenarios of reduction of the emission lead to a decrease in the SO₂ concentrations;
- the better results are demonstrated by the WAM 2030 scenario compared with the reference period 2005 and also compared with other scenarios;
- the bias of the SO₂ concentrations also has a negative sign, and the percentage of reduction is very well expressed and shows very good results for taking measures;
- the reduction reaches 50% almost over the whole territory of the country and for all scenarios;
- a reduction of 70% is achieved for some areas in the southwest part.

For all the results, we can conclude that more stringent measures lead to more effective results, whereas this study is about the WAM scenario.

ACKNOWLEDGEMENTS

This work has been carried out in the framework of the National Science Program "Environmental Protection and Reduction of Risks of Adverse Events and Natural Disasters", approved by the Resolution of the Council of Ministers № 577/17.08.2018 and has been accomplished with the financial support by the Ministry of Education and Science (MES) of Bulgaria (Agreement № D01-279/03.12.2021) and the project "NATIONAL GEO-INFORMATION CENTER", subject of the National Road Map for Scientific Infrastructure 2017–2023, funded by Contr. No D01-161/28.08.2018.

REFERENCES

- Byun, D., Young J., Gipson G., Godowitch J., Binkowski F.S., Roselle S., Benjey B., Pleim J., Ching J., Novak J., Coats C., Odman T., Hanna A., Alapaty K., Mathur R., McHenry J., Shankar U., Fine S., Xiu A., and Jang C., 1998: *Description of the Models-3 Community Multiscale Air Quality (CMAQ)*

- Modeling System*, 10th Joint Conference on the Applications of Air Pollution Meteorology with the A&WMA, 11-16 January 1998, Phoenix, Arizona, 264-268.
- Byun, D. and Ching, J., 1999: *Science Algorithms of the EPA Models-3 Community Multiscale Air Quality (CMAQ) Modeling System*. EPA Report 600/R-99/030, Washington, D.C., EPA/600/R-99/030 (NTIS PB2000-100561).
https://cfpub.epa.gov/si/si_public_record_report.cfm?Lab=NERL&dirEntryId=63400
- CEP, 2003: *Sparse Matrix Operator Kernel Emission (SMOKE) Modeling System*, University of Carolina, Carolina Environmental Programs, Research Triangle Park, North Carolina.
- CMAS Homepage, <http://www.cmaq-model.org>, last accessed 2022/05/10
- Denier van der Gon, H., Visschedijk, A., van de Brugh, H., Droge, R., 2010, *A high resolution European emission data base for the year 2005*, TNO-report TNO-034-UT-2010-01895 RPT-ML, Apeldoorn, The Netherlands (2010).
- Directive (EU) 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC <http://data.europa.eu/eli/dir/2016/2284/oj>
- EEA Technical report No 4/2015, *Projections in hindsight an assessment of past emission projections reported by Member States under EU air pollution and GHG legislation*
<https://www.eea.europa.eu/publications/projections-in-hindsight/download>
- Gadzhev, G., Ganey, K., Prodanova, M., Syrakov, D., Atanasov, E., Miloshev, N., 2013a, *Multi-scale Atmospheric Composition Modelling for Bulgaria*, NATO Science for Peace and Security Series C: Environmental Security, 137, 381-385.
- Gadzhev, G.K., Ganey, K.G., Prodanov, M., Syrakov, D.E., Miloshev, N.G., Georgiev, G.J., 2013b, *Some numerically studies of the atmospheric composition climate of Bulgaria*, AIP Conference Proceedings, 1561, 100-111. DOI: 10.1063/1.4827219 529
- Gadzhev, G., Ganey, K., Miloshev, N., Syrakov, D., Prodanova, M., 2013c, *Numerical Study of the Atmospheric Composition in Bulgaria*, Computers and Mathematics with Applications, 65, 402 – 422.
- Georgieva E., Hristova E., Syrakov D., Prodanova M., Batchvarova E., 2017, *Preliminary evaluation of CMAQ modelled wet deposition of sulphur and nitrogen over Bulgaria*. HARMO 2017 – 18th International Conference on Harmonisation within Atmospheric Dispersion Modelling for Regulatory Purposes, Proceedings 2017 – October, 51 – 55.
- Georgieva E., Syrakov D., Hristova E., Prodanova M., Gospodinov I., 2019, *Comparison of EMEP and WRF_CMAQ modelling results for deposition estimates in Bulgaria for 2016 and 2017*, 19th International Conference on Harmonisation within Atmospheric Dispersion Modelling for Regulatory Purposes, Harmo 2019.
- Georgieva, I., 2014, *Study of the air quality index climate for Bulgaria*, Proc. of the international conference on numerical methods for scientific computations and advanced applications, May 19 – 22, 2014, Bansko, ISBN978-954-91700-7-8, 39– 42.
- Shamarock, W.C., Joseph B., Dudhia J., Gill Dale M., Wang W., Jordan G., 2007: *A Description of the Advanced Research WRF Version 2*. <http://www.dtic.mil/dtic/tr/fulltext/u2/a487419.pdf>
- Syrakov, D., Etropolska, I., Prodanova, M., Ganey, K., Miloshev, N., Slavov, K., 2012, *Operational Pollution Forecast for the Region of Bulgaria*, American Institute of Physics, Conf. Proc. 1487, 88 - 94; DOI: 10.1063/1.4758945.
- Syrakov, D., Prodanova, M., Etropolska, I., Slavov, K., Ganey, K., Miloshev, N., Ljubenov T., 2014, *A multy-domain operational chemical weather forecast system*, Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics), 8353 LNCS, 413-420.
- Syrakov, D., Prodanova, M., Georgieva, E., Etropolska, I., Slavov, K., 2015a, *Impact of NOx emissions on air quality simulations with the Bulgarian WRF-CMAQ modelling system*, International Journal of Environment and Pollution, 57, 3-4, 285-296.
- Syrakov D., Georgieva E., Prodanova M., Hristova E., Gospodinov I., Slavov K., Veleva B., 2019, *Application of WRF-CMAQ model system for analysis of sulfur and nitrogen deposition over Bulgaria*, Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics), 11189 LNCS, 474 – 482.
- UCAR/ NCAR Homepage, <http://www.wrf-model.org>, last accessed 2022/05/10